

2

AD-A248 098

U
SECURITY



ITATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS														
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release Distribution unlimited														
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE																	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 8			5. MONITORING ORGANIZATION REPORT NUMBER(S)														
6a. NAME OF PERFORMING ORGANIZATION Brigham Young University		6b. OFFICE SYMBOL (If applicable) BYU		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research													
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Provo, UT 84602			7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Arlington, VA 22217-5000														
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable) ONR		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-91-J-1710													
8c. ADDRESS (City, State, and ZIP Code) 800 No. Quincy Street Arlington, VA 22217-5000			10. SOURCE OF FUNDING NUMBERS <table border="1"><tr><td>PROGRAM ELEMENT NO</td><td>PROJECT NO</td><td>TASK NO</td><td>WORK UNIT ACCESSION NO.</td></tr><tr><td></td><td></td><td></td><td></td></tr></table>			PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO.								
PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO.														
11. TITLE (Include Security Classification) Solid State Structures of Some Dithionoamido-18-crown-6 Ligands																	
12. PERSONAL AUTHOR(S) P. Huszthy, J.S. Bradshaw, N.K. Dalley and R.M. Izatt																	
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 1992, March 24													
15. PAGE COUNT																	
16. SUPPLEMENTARY NOTATION																	
17. COSATI CODES <table border="1"><tr><th>FIELD</th><th>GROUP</th><th>SUB-GROUP</th></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr></table>			FIELD	GROUP	SUB-GROUP										18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP															
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The structures of three Dithionoamido-18-crown-6 ligands have been determined by X-ray diffraction studies. The results show that the carbon and oxygen atoms on the macroring opposite the pyridine ring are disordered. A serious conformation change is noted when methyl substituents are bonded to the amide nitrogen atoms.																	
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS																	
21. ABSTRACT SECURITY CLASSIFICATION Unclassified																	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Harold Guard			22b. TELEPHONE (Include Area Code) (202) 696-4409		22c. OFFICE SYMBOL ONR												


92-07982



CONTINUED ON BACK

92 3 30 031

Figure 1. Chiral Diamido-, Dithionoamido-, Diaza- and Azapyridino-18-crown-6 Ligands



- | | |
|---|-----------------------------------|
| 1, X = NH; Y = O; R = benzyl (S,S) | |
| 2, X = NH; Y = S; R = benzyl (S,S) | 10, R = H (S,S) |
| 3, X = NH; Y = H ₂ ; R = benzyl (S,S) | 11, R = C(O)CH ₃ (S,S) |
| 4, X = NH; Y = O; R = phenyl (S,S) | |
| 5, X = NH; Y = S; R = phenyl (S,S) | |
| 6, X = NH; Y = H ₂ ; R = phenyl (S,S) | |
| 7, X = NCH ₃ ; Y = O; R = phenyl (S,S) | |
| 8, X = NCH ₃ ; Y = S; R = phenyl (S,S) | |
| 9, X = NCH ₃ ; Y = H ₂ ; R = phenyl (S,S) | |

The structures of 2, 5, and 8 were determined by X-ray diffraction studies. The discussion of the experimental procedures used in these studies and tables of structure determination summaries, atomic parameters, and torsion angles are included in the supplementary material. The molecule of 5 contained a 2-fold axis. The unit cells of both 5 and 8 contain two chemically similar but crystallographically distinct molecules of 5 and 8. Unfortunately, the number of single crystal data for each structure was not sufficient for a full anisotropic refinement, but by blocking atoms so that the number of parameters to be refined in each block was not so large, it was possible to refine most of the non-hydrogen atoms of the three structures anisotropically. Specifically, all non-hydrogen atoms of 2, all non-hydrogen atoms of one molecule of 5 and all non-

hydrogen atoms of the macrocyclic ring, pyridine ring, and the sulfur atom of both molecules of **8** were refined anisotropically. One molecule of **5**, which was badly disordered, and the methyl and phenyl carbon atoms of the two molecules of **8** were refined isotropically. The resulting *R* values were for **2**, *R* = 0.057, *R*_w = 0.0645; for **5**, *R* = 0.097, *R*_w = 0.085; and for **8**, *R* = 0.079, and *R*_w = 0.043. Details of the blocking of parameters and the treatment of hydrogen atoms of the three structures are contained in the supplementary material.

Computer drawings of the three molecules are shown in figures 2, 3 and 4. Only one of the two molecules of **5** and **8** were included as the conformations of the pairs are similar. These drawings clearly establish the structural formulas of the molecules, and also show the conformation of each molecule. In all three molecules, the 18-member ring is severely strained and deviates from the expected conformation of 18-crown-6 type molecules. This is established by the torsion angles listed in the supplementary material. This strain is expected because of the presence of the aromatic pyridine ring in the macrocyclic ring, and also the replacement of two of the oxygens of the ring by two nitrogens. As a result of these features, the carbon and oxygen atoms opposite the pyridine in the macrocyclic ring are disordered (see the thermal parameters for each structure in the supplementary materials). Perhaps the most interesting feature is the conformational change brought about by the presence of the methyl groups bonded to the nitrogens of the ether ring of **8**. This effect is shown in the deviations of the S and N atoms from the plane of the pyridine ring. The deviations of the nitrogen and sulfur atoms from the plane calculated for the pyridine and neighboring carbon atoms bonded to the pyridine are shown in Table II.




Figure 2. X-ray crystal structure of **2** drawn with SHELXTL-PLUS.³⁹ Hydrogen atoms were omitted for clarity.




Figure 3. X-ray crystal structure of the unprimed molecule of **5** drawn with SHELXTL-PLUS.³⁹ The primed molecule and hydrogen atoms were omitted for clarity.




Figure 4. X-ray crystal structure of the unprimed molecule of **8** drawn with SHELXTL-PLUS.³⁹ The primed molecule and hydrogen atoms were omitted for clarity.

Table II. Deviation of Nitrogen and Sulfur Atoms from the Plane of the Pyridine and Neighboring Carbon Atoms

Molecule	Deviation of S(Å)		Deviation of N(Å)	
2	S3	0.46	N4	-0.39
	S17	-0.13	N16	0.21
5	S3	-0.20	N4	0.20
	S3'	-0.01	N4'	0.12
8	S3	-1.56	N4	1.02
	S17	1.16	N14	-0.90
	S3'	1.48	N4'	-0.99
	S17'	-1.21	N16'	1.02